

The instruction on medicinal use of preparation Filtrum®-STI

Registration number: R No. 001189/01

Dosage form: pills

Contents: Active substance: hydrolyzed lignin – 0.4 g,

Pharmacotherapeutic group: enterosorbent

Pharmacological properties of Filtrum®-STI – it is a natural enterosorbent, which consists of the products of hydrolysis of wood components – lignin polymer, structural elements of which are phenylpropane and hydrocellulose. It has high sorption activity and non-specific detoxification action. It binds and discharges from the organism pathogenic bacteria, bacterial toxins, medications, poisons, salts of the heavy metals, alcohol, allergens, as well as some excess products of metabolism, including bilirubin, cholesterol, urea, metabolites responsible for the development of endogen toxemia. It is non-toxic, does not absorb, and is completely discharged from the intestine within 24 hours.

Indications for use

As detoxification medication for adults and children in cases of exogenous and endogenous intoxications of different origin:

- Acute intoxications with drugs, alkaloids, salts of the heavy metals, alcohol and other poisons;
- Complex treatment of food toxic infections, salmonellosis, dysentery, dyspepsia;
- Pyoinflammatory illnesses, accompanied with significant intoxication;
- Hyperbilirubinemia and hyperasotemia (liver and kidney insufficiency);
- Food and drug allergies;
- Prophylactics of chronic intoxications in workers of hazardous industries.

Contraindications

Individual intolerance to preparation. It is suggested not to use the drug in cases of stomach and duodenum ulcer illness exacerbations, intestinal atony.

Dosing and administration

Orally, preferably pre-crushed, with water one hour before the food intake or other drugs administration. The dose of the drug depends on age, body mass and severity of the illness, the drug is administered 3-4 times a day, medium one-time dose is for children: under 1 year old $\frac{1}{2}$ of the pill; 1–3 years old – $\frac{1}{2}$ -1 pill; 4–7 years old – 1 pill, 7–12 years old – 1–2 pills; for adults – 2-3 pills. The duration of treatment for acute conditions is 3-5 days, for allergic illnesses and chronic intoxications up to 14-21 days. Repeated courses of treatment after 2 weeks are upon doctor's advice.

Side effects

Rarely – allergic reactions, constipations. Long term administration of the drug may lead to disruption of vitamin and calcium absorption, so it is recommended to undertake prophylactic administration of multivitamins and calcium.

Special conditions

The drug may be used in complex therapy with other drugs, under condition of observing separate intake rule.

Sold in drugstores: Without receipt.

Manufacturer: AVVA RUS OJSC.